PUBLICATIONS:

1. **Madan I**, Harmanli OH, Dandolu V. Herpes in the Peripartum Period. Journal of Reproductive Medicine 2007; 52:539-540.

2. **Madan I**, Puri I, Jain NJ, Grotegut C, Nelson D, Dandolu V. Characteristics of Obstetric Intensive Care Unit Admissions in New Jersey. Journal of Maternal-Fetal and Neonatal Medicine 2009 Sep;22(9):785-90.

3. Mazaki-Tovi S, Vaisbuch E, Romero R, Kusanovic JP, Chaiworapongsa T, Kim SK, Nhan- Chang CL, Gomez R, Alpay Savasan Z, **Madan I**, Yoon BH, Yeo L, Mittal P, Ogge G, Gonzalez JM, Hassan SS. Maternal and neonatal circulating visfatin concentrations in patients with pre-eclampsia and a small-for-gestational age neonate. Journal of Maternal-Fetal and Neonatal Medicine 2010 Oct;23(10):1119-28

4. **Madan I**, Romero R, Kusanovic JP, Mittal P, Chaiworapongsa T, Dong Z, Mazaki- Tovi S, Vaisbuch E, Alpay-Savasan Z, Yeo L, Hassan SS. The frequency and clinical significance of intra-amniotic infection in women with placenta previa and vaginal bleeding: an unexpected observation. Journal of Perinatal Medicine 2010 May;38(3):275-9.

5. Alpay-Savasan Z, Romero R, Chaiworapongsa T, Kusanovic JP Kim SK, Romero R, Chaiworapongsa T, Kusanovic JP, Mazaki-Tovi S, Vaisbuch E, Mittal P, Ogge G, Alpay-Savasan Z, **Madan I**, Dong Z, Yeo L, Hassan SS. Evidence in support for a role of anti-angiogenic factors in preterm prelabor rupture of membranes (PPROM). Journal of Maternal-Fetal and Neonatal Medicine 2010 Aug;23(8):828-41.

6. Chaiworapongsa T, Kusanovic JP, Alpay Savasan Z, Mazaki-Tovi S, Kim SK, Vaisbuch E, Tarca AL, Mittal P, Ogge G, **Madan I**, Dong Z, Yeo L, Hassan SS, Romero R. Fetal death: a condition with dissociation in the concentrations of soluble vascular endothelial growth factor receptor-2 between the maternal and fetal compartments. Journal of Maternal-Fetal and Neonatal Medicine 2010 Sep;23(9):960-72

7. Chaiworapongsa T, Romero R, Kusanovic JP, Alpay-Savasan Z, Kim SK, Mazaki-Tovi S, Vaisbuch E, Ogge G, **Madan I**, Dong Z, Yeo L, Mittal P, Hassan SS. Unexplained fetal death is associated with increased concentrations of antiangiogenic factors in amniotic fluid. Journal of Maternal-Fetal and Neonatal Medicine 2010 Aug;23(8):794